

Asbestos Is Still With Us

Repeat Call for a Universal Ban

COLLEGIUM RAMAZZINI

All forms of asbestos are proven human carcinogens, causing malignant mesothelioma and a host of other types of cancers. No exposure to asbestos is without risk; there is no safe threshold of exposure to asbestos. When evidence of the carcinogenicity of asbestos became incontrovertible, a worldwide ban was called for on asbestos use, mining, and manufacturing. Asbestos is now banned in 52 countries. Nonetheless, many countries still use, import, and export asbestos and asbestos-containing products; many countries that have banned other forms of asbestos still permit the use of chrysotile asbestos. This exemption has no basis in medical science, but reflects the political and economic influence of the asbestos industry. To protect the health of all people, the Collegium Ramazzini calls again on all countries of the world to join in the international endeavor to ban all forms of asbestos. An international ban on asbestos is urgently needed. *Key words:* asbestos; malignant mesothelioma; lung cancer; laryngeal cancer; ovarian cancer; cancer; asbestos mining; asbestos manufacturing; asbestos ban; chrysotile asbestos.

INT J OCCUP ENVIRON HEALTH 2010;16:351-355

INTRODUCTION

Asbestos is a term applied to six naturally occurring fibrous minerals. These minerals occur in two configurations: serpentine and amphibole. The only type of asbestos derived from serpentine minerals, chrysotile, also known as white asbestos, accounts for 95 percent of the asbestos ever used around the world, and it is the

only type of asbestos in commercial use today. Amphibole minerals include five asbestos species: amosite, crocidolite, tremolite, anthophyllite, and actinolite. The two forms of amphibole asbestos that previously were most commercially important—amosite, or brown asbestos, and crocidolite, or blue asbestos—are no longer in use.

Asbestos fibers can withstand fire, heat, and acid. They have great tensile strength. They provide thermal insulation and acoustic insulation. For these reasons, asbestos came into wide commercial use and gave rise to a burgeoning industry many years before its detrimental health effects, which often take years to appear, became known.

All forms of asbestos cause asbestosis, a progressive, debilitating fibrotic disease of the lungs. All forms of asbestos cause human cancer. All forms of asbestos cause malignant mesothelioma, lung, laryngeal, and ovarian cancers. All forms of asbestos may cause gastrointestinal and other cancers.¹

Asbestos was declared a proven human carcinogen by the US Environmental Protection Agency (EPA), the International Agency for Research on Cancer (IARC) of the World Health Organization (WHO), and the US National Toxicology Program (NTP) more than 20 years ago.²⁻⁴ The scientific community is in overwhelming agreement that there is no safe level of exposure to asbestos.⁵ Moreover, there is no evidence of a threshold level below which there is no risk of mesothelioma.⁶

THE ASBESTOS CANCER PANDEMIC

Occupational Exposures to Asbestos

About 125 million people around the world are exposed to asbestos in their work environments,⁷ and many millions more workers have been exposed to asbestos in years past. About 20 to 40 percent of adult men report having worked in occupations that may have entailed asbestos exposures.⁸ In the most highly affected age groups, mesothelioma may account for over 1% of all deaths.^{9,10} In addition to mesothelioma, five to seven percent of all lung cancers are potentially attributable to occupational exposures to asbestos.¹¹

Received from: Collegium Ramazzini, International Headquarters, Castello dei Pio, 41012 Carpi/Modena, Italy. The Collegium Ramazzini, an international academic society that examines critical issues in occupational and environmental medicine, is dedicated to the prevention of disease and the promotion of health. The Collegium derives its name from Bernardino Ramazzini, the father of occupational medicine, a professor of medicine of the Universities of Modena and Padua in the early 1700s. Currently, 180 renowned clinicians and scientists from around the world, each of whom has been elected to membership, comprise the Collegium. It is independent of commercial interests. Send correspondence to: General Secretariat, Castello di Bentivoglio, 40010 Bentivoglio, Bologna, Italy, 39 051 6640650; email: <collegium@ramazzini.it>.

Disclosures: The authors declare no conflicts of interest.

Worldwide, the yearly number of asbestos-related cancer deaths in workers is estimated to be 100,000 to 140,000. In Western Europe, North America, Japan, and Australia, 20,000 new cases of lung cancer and 10,000 cases of mesothelioma result every year from exposures to asbestos.¹² In the United Kingdom at least 3500 people die from asbestos-related illnesses each year, and this number is expected to increase to 5000 in future years.⁷ The British mesothelioma death-rate is now the highest in the world, with 1740 deaths in men (1 in 40 of all male cancer deaths below age 80) and 316 in women in 2006. About 1 in 170 of all British men born in the 1940s will die of mesothelioma.¹⁰ Australia's high incidence of mesothelioma is expected to reach 18,000 by 2020, with 11,000 cases yet to appear.¹³

The US National Institute for Occupational Safety and Health (NIOSH) estimates that current occupational exposures to asbestos even at the US Occupational Safety and Health Administration's (OSHA's) permissible exposure limit will cause five deaths from lung cancer and two deaths from asbestosis in every 1000 workers exposed for a working lifetime.¹⁴

Environmental Exposures to Asbestos

Non-occupational, environmental exposure to asbestos from the use of asbestos in construction materials is a serious and often neglected problem in countries throughout the world. In developed countries large quantities of asbestos are found today as a legacy of past construction practices in many thousands of schools, homes, and commercial buildings. And in developing countries, where asbestos continues to be used in large quantities in construction, asbestos-contaminated dust is now accumulating in thousands of communities.

More than 90 percent of the asbestos used worldwide today is used in the manufacture of asbestos-cement sheets and pipes, and most of this material is used in developing countries. Use of asbestos in these materials continues despite repeated warnings that the use of asbestos in these products is highly dangerous because of the large numbers of people exposed to the airborne dust and the extreme difficulty of controlling exposures once these materials have been disseminated into communities where people of all ages, including young children, are at risk of exposure.¹⁵ A pervasive problem with use of asbestos-containing materials in construction is that asbestos fibers are released to air and dust as these materials weather, erode, break, or are cut by saws and other power tools.⁷ Community-wide exposure to persons of all ages is the end result.

Both community-based and industrial exposures to asbestos and asbestiform fibers increase risks for mesothelioma.¹⁶ Thus a study of women residing in Canadian asbestos mining communities found a seven-fold increase in the mortality rate from pleural

cancer.¹⁷ The risk of developing asbestos-related cancer following in-home exposures in communities near Canadian mines over a 30-year period is estimated to be one in 10,000.¹⁸ Likewise, environmental exposures to asbestos waste on the surfaces of roads and yards in a contaminated community of 130,000 residents in The Netherlands result each year in several cases of malignant mesothelioma.¹⁹ And in a third example, the currently observed increase in female cases of mesothelioma in the United Kingdom, many with no occupational exposure to asbestos, suggests widespread environmental contamination.¹⁰

Chrysotile Asbestos

Chrysotile represents 95 percent of all the asbestos ever used worldwide. It is the only variety in international trade in the twenty-first century. There is general agreement among scientists and physicians, and widespread support from numerous national health agencies in countries around the world, United Nations agencies, and the World Trade Organization, that chrysotile asbestos causes various cancers, including mesothelioma and lung cancer.²⁰⁻²⁹

Early suggestions that chrysotile might be less dangerous than other forms of asbestos have not been substantiated. Although chrysotile accounts for almost all the asbestos ever used, the asbestos industry continues to claim that asbestos-related cancers are the result of the amphibole varieties.^{30,31} Defenders of the chrysotile asbestos industry contend—falsely, and despite all of the abundant medical and scientific evidence to the contrary—that, “Exposure to chrysotile in a pure form seems likely to present a very low if any risk of mesothelioma.”³²

The Chrysotile Institute, a registered lobbying group for the Quebec asbestos mining industry, takes the position that chrysotile can be handled safely.³³ But refuting this scientifically untenable and highly misleading position are numerous epidemiologic studies, case reports, controlled animal experiments, and toxicological studies which show clearly and consistently that chrysotile is highly dangerous and that it is fully capable of causing cancer.^{12,34-39} These studies demonstrate that the so-called “controlled use” of asbestos is a fallacy.⁴⁰ Workers exposed to chrysotile fiber alone have excessive risks of lung cancer and excessive deaths from mesothelioma.⁴¹⁻⁴³

The Canadian Medical Association, the Canadian Cancer Society, and Canada's leading health experts oppose the export of asbestos to developing countries. The National Public Health Institute of Quebec (INSPQ) has published fifteen reports, all of them showing a failure to achieve “controlled use” of asbestos in Quebec itself. Pat Martin, a member of Canada's parliament and former asbestos miner asks, “If we in the developed world haven't found a way to

handle chrysotile safely, how can we expect them to do so in developing nations?"⁴⁴

CURRENT PRODUCTION AND USE OF ASBESTOS

Despite all that is known about the health effects of asbestos, annual world production remains at over two million tons. This level of production has remained steady following a 50% decline in the 1990s. Russia is now the leading producer of asbestos worldwide, followed by China, Kazakhstan, Brazil, Canada, Zimbabwe, and Colombia. These six countries accounted for 96% of the world production of asbestos in 2007.⁴⁵ Russia has mines rich enough in asbestos deposits to last for more than 100 years at current levels of production. The majority of the 925,000 tons of asbestos extracted annually in Russia is exported.

Asbestos is now banned in 52 countries,⁴⁶ including all EU member countries, and safer products have replaced many that were once made with asbestos. Virtually all of the polymeric and cellulose fibers used instead of asbestos in fiber-cement sheets are greater than 10 microns in diameter and hence are non-respirable. Nonetheless, these 52 countries make up less than a third of WHO member countries.

Unfortunately, a much larger number of WHO member countries still use, import, and export asbestos and asbestos-containing products.²⁸ These are mostly developing countries, and over 70 percent of the world production of asbestos is used today in Asia and Eastern Europe, in countries desperate for industrial growth and often naïve to the health effects of occupational and environmental exposures to asbestos. A recent article in *The Lancet* notes that, "Vast development projects in Asia are largely responsible for maintaining the [chrysotile asbestos] market. In particular, India's asbestos industry is burgeoning."⁴⁴

In many countries that have banned other forms of asbestos, the "controlled use" of chrysotile asbestos is still permitted, despite all medical and scientific information to the contrary. This exemption reflects the size of the asbestos industry, its pervasive influence, and the importance of asbestos mining and manufacture to the economy. The toll in most countries still using large amounts of asbestos may never be fully recorded.

In developing countries, where too often there exists little or no protection of workers and communities, the asbestos cancer pandemic may be the most devastating. China is by far the largest consumer of asbestos in the world today, followed by India, Russia, Kazakhstan, Thailand, Ukraine, and Uzbekistan.

Position of United Nations Agencies on Asbestos

International organizations have condemned the continuing use of chrysotile asbestos.⁴⁷ In 2006, the WHO

called for the elimination of diseases associated with asbestos.²⁸ The WHO supports individual countries in developing national plans to ban asbestos and eliminate asbestos disease. The WHO has stated that "the most efficient way to eliminate asbestos-related disease is to stop using all types of asbestos."

The International Labour Organization (ILO) has expressed concern about an evolving epidemic of asbestos-related diseases, and has passed a resolution to promote a worldwide asbestos ban.²²

The World Trade Organization has accepted the conclusion that the so-called "controlled use" of asbestos is a fallacy.⁴⁸

The Rotterdam Convention is an international treaty intended to regulate global trade in dangerous chemicals—chemicals that have been banned or severely restricted because of their hazards to human health or the environment. It was enacted in 2004, and 131 nations are currently parties to the Convention. The goal is to protect the world's most vulnerable countries—developing countries and countries with economies in transition—against importation without their prior knowledge or consent of hazardous pesticides and other regulated chemicals.

Prior informed consent (PIC) is the core principle of the Rotterdam Convention. This legally binding procedure requires that governments in all countries be provided full information prior to importation about the risks to health and the environment of each of the hazardous materials regulated by the Convention. Annex III of the Rotterdam Convention contains a list of the chemicals—37 in number—currently regulated by the Convention.

Repeated efforts to include chrysotile asbestos under the Rotterdam Convention have failed because of the Convention's requirement for unanimity and the determined opposition of asbestos mining and manufacturing countries.⁴⁹ At the 2008 conference of parties on the Convention, opposition to chrysotile asbestos was led by Russia, India, and Kazakhstan. A few asbestos importing countries thwarted the will of over 100 other countries.

CONCLUSION: THE NEED FOR A UNIVERSAL BAN ON ASBESTOS

The profound tragedy of the asbestos pandemic is that virtually all illnesses and deaths related to asbestos are preventable. Safer substitutes for asbestos exist, and they have been introduced successfully in many nations. Asbestos-cement (A-C) pipes, sheets, and water storage tanks account for 90 percent of asbestos used in the world today. Substitutes for A-C water pipe include ductile iron pipe, high-density polyethylene pipe, and metal-wire-reinforced concrete pipe. Many substitutes exist for roofing, interior building walls, and ceilings, including fiber-cement flat and corrugated sheet products, made with

polyvinyl alcohol fibers and cellulose fibers. For roofing, lightweight concrete tiles can be made and used in the most remote locations, using locally available plant fibers including jute, hemp, sisal, palm nut, coconut coir, and wood pulp. Galvanized iron roofing and clay tiles are among the other alternative materials.⁵⁰

If global use of asbestos were to cease today, a decrease in the incidence of asbestos-related diseases would become evident only two or more decades from now.²⁸ This delay is a consequence of the long latency period associated with the diseases caused by asbestos. In the case of mesothelioma, the latency between exposure and disease may be as long as 40 to 50 years.

The asbestos cancer pandemic may take as many as 10 million lives before asbestos is banned worldwide and all exposure is brought to an end.^{47,51} In this conservative estimate, it is assumed that asbestos exposures are going to cease and that the epidemic will run itself out. But in fact, the world's current production of asbestos continues at an alarming rate, and therefore these figures may be underestimates of the true reality of this pandemic.

An international ban on the mining and use of asbestos is urgently needed. The risks of exposure to asbestos cannot be controlled by technology or by regulation of work practices. Scientists and responsible authorities in countries allowing the use of asbestos should have no illusion that "controlled use" of chrysotile asbestos is an effective alternative to a ban on all use of asbestos.^{52,53} Even the best workplace controls cannot prevent occupational and environmental exposures to products in use or to waste. Safer substitute products are available and in use in countries all over the world where asbestos is banned.

To protect the health of all people in the world—industrial and construction workers, as well as their families—now and in future generations, the Collegium Ramazzini calls again today on all countries of the world, as we have repeatedly in the past, to join in the international endeavor to ban all forms of asbestos. An international ban on asbestos is urgently needed.

References

1. Straif K, Benbrahim-Tallaa L, Baan R, Grosse Y, Secretan B, El Ghissassi F, Bouvard V, Guha N, Freeman C, Galichet L, Coglianò V. [WHO International Agency for Research on Cancer Monograph Working Group]. A review of human carcinogens—Part C: Metals, arsenic, dusts, and fibres. *Lancet Oncol.* 2009;10(5):453-454.
2. United States Environmental Protection Agency (EPA). Airborne asbestos health assessment update. EPA/6000/8-84/003E. Washington, DC: EPA; 1986.
3. International Agency for Research on Cancer (IARC). Asbestos: Monograph on the evaluation of carcinogenic risk to man. Vol 14. Lyon, France: IARC; 1977.
4. National Toxicology Program (NTP). Report on carcinogens. 1st ed. Washington, DC: US Department of Health and Human Services, Public Health Service; 1980.
5. Welch LS. Asbestos exposure causes mesothelioma, but not this asbestos exposure: An amicus brief to the Michigan Supreme Court. *Int J Occup Environ Health.* 2007;13:318-327.
6. Hillerdal G. Mesothelioma: Cases associated with non-occupational and low dose exposures. *Occup Environ Med.* 1999; 56(8):505-513.
7. Egilman D, Fehnel C, Bohme SR. Exposing the "myth" of ABC, "anything but chrysotile": A critique of the Canadian asbestos mining industry and McGill University chrysotile studies. *Am J Ind Med.* 2003;44:540-557.
8. Goldberg M, Banaei A, Goldberg S, Anvert B, Luce D, Gueguen A. Past occupational exposure to asbestos among men in France. *Scand J Work Environ Health.* 2000;26:52-61.
9. Driscoll T, Nelson DI, Steenland K, Leigh J, Concha-Barrientos M, Fingerhut M. The global burden of disease due to occupational carcinogens. *Am J Ind Med.* 2005;48(6):419-431.
10. Rake C, Gilham C, Hatch J, Darnton A, Hodgson J, Peto J. Occupational, domestic and environmental mesothelioma risks in the British population: A case-control study. *Br J Cancer.* 2009;100(7):1175-1183.
11. Tossavainen A. Global use of asbestos and incidence of mesothelioma. *Int J Occup Environ Health.* 2004;10:22-25.
12. Tossavainen A. Asbestos, asbestosis and cancer: The Helsinki criteria for diagnosis and attribution. Consensus report. *Scand J Work Environ Health.* 1997;23:311-316.
13. Leigh J, Driscoll T. Malignant mesothelioma in Australia, 1945-2002. *Int J Occup Environ Health.* 2003;9(3):206-217.
14. Stayner L, Smith R, Bailer J, Gilbert S, Steenland K, Dement J, Brown D, Lemen R. Exposure-response analysis of risk of respiratory disease associated with occupational exposure to chrysotile asbestos. *Occ Env Med.* 1997;54:646-652.
15. World Health Organization (WHO). Environmental health criteria: 203: Chrysotile asbestos. Geneva, Switzerland: WHO; 1998.
16. Pasetto R, Comba P, Marconi A. Mesothelioma associated with environmental exposures. *Med Lav.* 2005;96(4):330-337.
17. Camus M, Siemiatycki J, Meek B. Nonoccupational exposure to chrysotile asbestos and the risk of lung cancer. *N. Engl. J. Med.* 1998;338: 1565-71.
18. Marier M, Charney W, Rousseau R, Lanthier R, Van Raalte J. Exploratory sampling of asbestos in residences near Thetford Mines: The public health threat in Quebec. *Int J Occup Environ Health.* 2007;13:386-397.
19. Driece HA, Siesling S, Swuste PH, Burdorf A. Assessment of cancer risks due to environmental exposure to asbestos. *J Expo Sci Environ Epidemiol.* 2009; Oct 28. [Epub ahead of print].
20. American Conference of Governmental Industrial Hygienists (ACGIH). Asbestos: TLV chemicals. 7th ed. Publication #7DOC-040. Cincinnati, OH: ACGIH; 2001.
21. Agency for Toxic Substances and Disease Registry (ATSDR). Toxicological profile for asbestos. US Department of Health and Human Services: Atlanta, GA; 2001.
22. International Labour Organization (ILO). ILO adopts new measures on occupational safety and health, the employment relationship, asbestos. [Internet] 2006. Geneva, Switzerland: ILO. Available from: <http://www.ilo.org/public/english/bureau/inf/pr/2006/34.htm>.
23. International Social Security Association (ISSA). Declaration on asbestos Beijing 2004. The Special Commission on Prevention of the International Social Security Association (ISSA), assembled in Beijing on 16 September 2004 on the occasion of the 28th General Assembly of the Association, issues this appeal to governments in asbestos-producing countries. Available at: <http://www.issa.int/aiss/Resources/Resources/Declaration-on-Asbestos-Beijing-2004>.
24. National Toxicology Program (NTP). Report on carcinogens. 11th ed. Washington, DC: US Department of Health and Human Services, Public Health Service; 2004.
25. National Cancer Institute (NCI). Factsheet—Asbestos: Questions and answers. Bethesda, MD: National Institutes of Health; 2003.
26. Occupational Safety and Health Administration (OSHA). Occupational exposure to asbestos: Final rule. *Fed Reg.* 1994; 59:40964-41162.
27. United Nations Environment Program (UNEP), International Labor Office (ILO), World Health Organization (WHO). Chrysotile asbestos, environmental health criteria 203. Geneva, Switzerland: WHO; 1998.
28. World Health Organization (WHO). Elimination of asbestos related diseases. Geneva, Switzerland: WHO; 2006.

29. World Trade Organization (WTO). European communities—Measures affecting asbestos and asbestos-containing products. Report of the Panel. WT/DS135/R. 2000 Sept 18. Available at: <http://docsonline.wto.org/DDFDocuments/t/WT/DS/135R-00.doc>.
30. McCulloch J. Saving the asbestos industry, 1960 to 2006. Public Health Reports. 2006;121:609-614.
31. Renner R. Chrysotile on ice. Environ Health Perspect. 2007; 115:3:130.
32. Gibbs GW, Berry G. Mesothelioma and asbestos. Regul Toxicol Pharmacol. 2008;52(1 Suppl):S223-31.
33. Chrysotile Institute. The crusade against chrysotile must end. Newsletter [Internet]. 2008;7(2):2. Available from: http://www.chrysotile.com/data/newsletter/Chrysotile_Dec2008_EN.pdf.
34. Bang KM, Pinheiro GA, Wood JM, Syamlal G. Malignant mesothelioma mortality in the United States, 1999-2001. Int J Occup Environ Health. 2006;12(1):2-15.
35. Landrigan P, Nicholson WJ, Suzuki Y, LaDou J. The hazards of chrysotile asbestos: A critical review. Ind Health. 1999;37:271-280.
36. Lemen RA. Chrysotile asbestos as a cause of mesothelioma: Application of the Hill causation model. Int J Occ Env Health. 2004a;10:233-239.
37. Lin RT, Takahashi K, Karjalainen A, Wilson D, Kameda T, Chan CC, Wen CP, Furuva S, Higashi T, Chan LC, Ohtaki M. Ecological association between asbestos-related diseases and historical asbestos consumption: An international analysis. Lancet. 2007; 20(9564):844-849.
38. Smith AH, Wright CC. Chrysotile asbestos is the main cause of pleural mesothelioma. Am J Ind Med. 2006;30:252-266.
39. Stayner LT, Dankovic DA, Lemen RA. Occupational exposure to chrysotile asbestos and cancer risk: A review of the amphibole hypothesis. Am J Pub Health. 1996;86:179-186.
40. Lemen RA. Asbestos in brakes: Exposure and risk of disease. Am J Ind Med. 2004b;45(3):229-237.
41. Frank AL, Dodson RF, Williams MG. Carcinogenic implications of the lack of tremolite in UICC reference chrysotile. Am. J. Ind. Med. 1998;34:314-317.
42. Li L, Sun TD, Zhang X, Lai RN, Li SY, Fan XJ, Morinaga K. Cohort studies on cancer mortality among workers exposed only to chrysotile asbestos: A meta-analysis. Biomed Environ Sci. 2004;17(4):459-468.
43. Mirabelli D, Calisti R, Barone-Adesi F, Fornero E, Merletti F, Magnani C. Excess of mesotheliomas after exposure to chrysotile in Balangero, Italy. Occup Environ Med. 2008;65(12):815-819.
44. Burki T. Health experts concerned over India's asbestos industry. [Internet]. The Lancet. 2010;375(9715):626-627. Available from: [http://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(10\)60251-6/fulltext](http://www.thelancet.com/journals/lancet/article/PIIS0140-6736(10)60251-6/fulltext).
45. United States Geological Survey (USGS). 2008 minerals yearbook. Volume I. Metals and minerals. Asbestos. [Internet]. Washington, DC: US Department of the Interior; 2008. Available from: <http://minerals.usgs.gov/minerals/pubs/commodity/asbestos/index.html#myb>
46. International Ban Asbestos Secretariat. (IBAS). [Internet] 2010 [cited 2010 Mar 8]. Available from: http://ibasecretariat.org/alpha_ban_list.php.
47. LaDou J. The asbestos cancer epidemic. Env Health Perspect. 2004;112:285-290.
48. Castleman B. WTO confidential: The case of asbestos. Int J Health Serv. 2002;32:489-501.
49. Terracini B. Rotterdam Convention: Chrysotile is still in the waiting list. Epidemiol Prev. 2008;32(6):275-276. (Italian)
50. World Bank Group (WBG). Asbestos: Occupational and community health issues. Good Practice Note. [Internet] 2009. World Bank Operations Policy and Country Services. Available from: <http://siteresources.worldbank.org/EXTPOPS/Resources/AsbestosGuidanceNoteFinal.pdf>
51. Leigh J. Asbestos-related diseases: International estimates of future liability. Abstract. 5th International Congress on Work Injuries Prevention, Rehabilitation and Compensation & 2nd Australian National Workers Compensation Symposium. Adelaide, Australia; 18-21 March, 2001.
52. Castleman B. 2003. "Controlled use" of asbestos. Int J Occup Environ Health. 2003;9:294-298.
53. Egilman D, Roberts M. Controlled use of asbestos. Int J Occup Environ Health. 2004;10:99-103.